

Parton content of hadron interactions at Tevatron and LHC

I.M. Dremin, V.A. Nechitailo

Lebedev Physical Institute, Moscow

Vietnam, December 2011

The independent pair parton interactions (IPPI) model:


1. High energy particles are "clouds" of partons.
2. Each pair of colliding partons is independent of others and creates new particles according to NBD-distribution of multiplicity.

The main equation:

$$P(n; m, k) = \sum_{j=1}^{j_{max}} w_j P_{NBD}(n; jm, jk). \quad (1)$$

$P(n; m, k)$ is the probability to create n particles, m and k are the parameters of the NBD-distribution, w_j is the probability for j parton pairs to be active at a given energy, $\sum_{j=1}^{j_{max}} w_j = \sum_{j=1}^{j_{max}} w_1^j = 1$, j_{max} is the maximum number of the active parton pairs.

- Experimental indications: single NBD fits at energies up to 200 GeV, then the distributions widen.
- Interpretation: 1 pair of partons is active and leads to NBD at lower energies while their number increases with energy.



The decomposition of the multiplicity distribution at Tevatron $\sqrt{s}=1.8$ TeV into 1, 2, 3 and 4 parton-parton interactions

How one gets the main equation.

Main property:

The two folded NBD lead to NBD!


$$P(n; m, k) = \sum_{j=1}^{j_{max}} w_j P_j(n; m, k) = \sum_{j=1}^{j_{max}} w_j \sum_{(n_p)} \prod_{p=1}^j P_{NBD}(n_p; m, k). \quad (2)$$

n_p is the number of particles created by the p th pair,
 $\sum_{(n_p)}$ denotes the folder of NBD distributions with the sum of those parton interactions where $\sum_{p=1}^j n_p = n$.


The summation in Eq. (2) gives rise to the main equation.

NBD-reminder:


$$P_{NBD}(n; m, k) = \frac{\Gamma(n+k)}{\Gamma(n+1)\Gamma(k)} \left(\frac{m}{k}\right)^n \left(1 + \frac{m}{k}\right)^{-n-k}$$



The fit by the [IPPI-model](#) (dotted) of the multiplicity distribution at 1.8 TeV.



The fits by the **IPPI**-model (line) and by the **QGSM**-model (dots) of the multiplicity distribution at 7 TeV ($|\eta| < 2.4$, CMS-data).



The values of the parameter m (i.e. of the effective average multiplicity for a single parton pair) at different energies and rapidity windows.

The number of active pairs (j_{max}) is shown near each point.

IMPORTANT TECHNICALITIES!

The requirement of independence of m on the ranks of moments of the distribution imposes restrictions on the parameter k .

We use the properties of factorial (F_q) and cumulant (K_q) moments of the multiplicity distributions (as well as of their ratio H_q) in fits of experimental data to show how well this requirement is fulfilled.

For more details see the papers in Phys. Rev. or arXiv:hep-ph.


$$\begin{aligned} F_q &= \sum_n P(n) n(n-1)\dots(n-q+1) \\ &= \sum_{j=1}^{j_{\max}} w_j \frac{\Gamma(jk+q)}{\Gamma(jk)} \left(\frac{m}{k}\right)^q = f_q(k) \left(\frac{m}{k}\right)^q \end{aligned} \quad (3)$$

where

$$f_q(k) = \sum_{j=1}^{j_{\max}} w_j \frac{\Gamma(jk+q)}{\Gamma(jk)} = k \sum_{j=1}^{j_{\max}} w_j j(jk+1)\dots(jk+q-1). \quad (4)$$

The selfconsistency of the IPPI-model asks for q -independence of m :

$$m = k \left(\frac{F_q^{exp}}{f_q(k)} \right)^{1/q} = \text{const.} \quad (5)$$



- The IPPI-model is proposed.
- The pairs of partons from colliding "clouds" are independent and each pair creates particles according to NBD-distribution.
- Experimental distributions are well described at different energies and rapidity windows with only two adjustable parameters m and j_{max} .
- The average multiplicity in collision of a single pair m and the number of active pairs j_{max} increase with energy logarithmically.
- The density of the parton medium increases with energy and asks for account of SOFT (not only HARD) multiparton interactions.
- The predictions at higher energies 14 and 100 TeV have been done.